

SAN JOSE GASTROENTEROLOGY

Phone (408) 347-9001 | Fax (408) 347-9004 | www.sjgi.com | contact@sjgi.com

INSTRUCTIONS FOR COLONOSCOPY BOWEL PREPARATION with PLENVU (SAME-DAY PREP)

For the best possible examination, please follow the instructions below carefully.

**** ONE (1) WEEK BEFORE THE COLONOSCOPY:** *For the best possible examination, please follow the instructions below carefully:*


- STOP iron pills, vitamins, herbs and supplements. AVOID vegetables, seeds and fibers (berries, multigrain bread, popcorn)
- **STOP ALL** medications that thin the blood (coumadin or warfarin, plavix, effient). It is important that you contact your primary physician to get approval when stopping these medications. You do not need to stop baby aspirin.
- **OBTAIN PLENVU** from your pharmacy. If this medication is replaced due to insurance requirement, please inform SJGI so new instruction can be given to you.

****ONE (1) DAY BEFORE THE COLONOSCOPY:**

- **CONTINUE** to take all diabetes medications regardless of taken time in morning or evening daily.
- **REDUCE** morning and evening doses of long acting insulin by one half.
- You will be on a CLEAR LIQUID DIET FOR THE ENTIRE DAY. **Do not fast or starve yourself. DO NOT eat any solid food for a full 24 hours prior to your colonoscopy.** It's important to stay hydrated before, during, and after taking PLENVU®.

For a full 24 hours prior to the colonoscopy, you must be on a clear liquid diet limited to water and the following:		DO NOT eat solid food DO NOT chew gum DO NOT drink any of the following: <ul style="list-style-type: none">▪ NO red or purple liquid▪ NO liquids that you can't see through▪ NO alcoholic beverages▪ NO milk products
Black coffee (no milk / no cream) Tea (black, green) Gatorade (clear) Popsicle w/o fruit or seed	7-UP Clear broth Apple Juice Jello (No red or purple)	

**** ON THE DAY OF THE COLONOSCOPY: TAKE DOSE 1 AT 6:00 AM**



- Use the mixing container to mix the contents of the Dose 1 pouch with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2 to 3 minutes. Take your time—slowly drink and finish the dose within 30 minutes.
- Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes.

Wait at least 2 hours from the start of dose 1, then take the 2nd dose



- Use the mixing container to mix the contents of Dose 2 (Pouch A and Pouch B) with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2 to 3 minutes. Take your time—slowly drink and finish the dose within 30 minutes.
- Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes.

- **STOP** all diabetes medications before procedure time. **OK** to resume all medications after procedure.
- **NOTE:** If you have nausea or vomit, **DO NOT STOP** taking the laxative. Rest for 30 minutes and then resume the preparation as soon as the nausea improves.
- **STOP DRINKING** liquids at least 2 hours before your colonoscopy.

PLEASE BE AWARE THAT:

- You must have a responsible adult (age 18 or older) to drive you home after your procedure. This does not include rideshare such as taxi, Uber. The sedative medication makes it unsafe to drive on the day of the procedure. **If you do not have a chaperone to drive you home, the appointment may need to be cancelled.**
- **You should not work after the colonoscopy. Plan to take the day off from work.**
- **DO NOT** bring children to your appointment. Leave jewelry and valuable items at home.



BRING all of your daily medication to the colonoscopy

TAKE your heart/blood pressure medication with *small sips of water before you come to your procedure*



If your colon output is **NOT** a clear yellow color or similar to water color after the laxative cleansing, inform the nurse when you arrive for your procedure.

If you have any questions, please contact our office at 408-347-9001. If you need to reschedule or cancel your appointment, please contact us at least **3 business days** in advance before your procedure.

Thank you very much for your cooperation.